

Credit: AFP/Getty Images

## Overview of Climate Hazards Relevant to Older Adults


Heidi Cullen - Climate Central July 19, 2011

#### Outline


- Variability & change within the physical climate system
- Ongoing & projected future climate impacts/threats
- Climate change and variability that posing significant risks for older adults



Jackson Pollock Number 8








Natural v. Man-made

Low-frequency:

Milankovitch cycles Ice sheets GHG's (like CO<sub>2</sub>)

----[High-frequency:

Solar variability (sunspots)
Volcanic eruptions
El-Niño
Us (fossil fuel burning, deforestation)





El Niño/La Niña: periodic warming cooling of tropical Pacific Ocean (2-7 years)

PNA: changes in atmospheric pressure between the Aleutian Low and the high pressure over the Rocky Mountains. Strongly influenced by El Nino.

NAO/AO: A large scale seesaw in atmospheric mass between the Azores High and Icelandic Low. Varies from year to year, but also can remain in one phase for several years.


www.climate.gov






El Niño: stronger winter storms in California and southern US with floods/landslides.

La Niña: drier than normal conditions in US Southwest in late summer through subsequent winter. Drier than normal conditions typically occur in Central Plains in fall and in the Southeast in winter. In contrast, the Pacific Northwest is more likely to be wetter than normal in the late fall and early winter. Stronger hurricane season.



Positive NAO: The eastern US experiences mild and wet winter conditions.

Negative NAO: The US east coast experiences more cold air outbreaks and hence snowy weather conditions.





Positive PNA: Colder temps in central/eastern US. Warmer temps in western US.

Negative PNA: Colder/wetter across the western US. Florida and southern parts of southeast drier/warmer in winter months.

the recent "Western ridge"/Positive PMA pattern

# Global warming is a trend.






David Keeling



Mauna Loa Observatory





## Climate Scientists Agree That...

- 1. Observed warming is unequivocal, human-caused and irreversible.
- 2. Natural climate variability is large.
- 3. Weather extremes are increasing.
- 4. The more CO<sub>2</sub> is emitted, the warmer it will become.
- 5. Emissions are currently above all projections.
- 6. Future climate scenarios are fundamentally about our choices.

Climate is what you expect, weather is what you get.

Mark Twain





Europe 2003

Summer of 2003 was probably the hottest in Europe since at latest AD 1500

Paris temperatures: 104F

Estimated 50,000 deaths


#### 2003 European Heat Wave



human influence at least doubled the chance of summers as hot as the one Europe saw in 2003

models predict that by 2040, the 2003-type summers will be happening every other year


#### Weather related mortality




 a) Location of Hurricane Landfalls, 1995 to 2000




 Percentage of Population Aged 65 or Older



 b) Location of Extreme Heat Events, 1995 to 2000



d) West Nile Virus Cases 2004




CCSP SAP 4.6163

heat waves: more frequent, extreme and long lasting


in same weather, more people die earlier in the summer than later in the summer

bodies adapt physiologically to producing more sweat glands over the span of a few weeks


little excess mortality during first 3 days, by fifth day excess deaths in dry heat are quadruple the number in humid heat (Philly)



The number of days in which the temperature exceeds 100°F by late this century, compared to the 1960s and 1970s, is projected to increase strongly across the United States. For example, parts of Texas that recently experienced about 10 to 20 days per year over 100°F are expected to experience more than 100 days per year in which the temperature exceeds 100°F by the end of the century under the higher emissions scenario.<sup>91</sup>




## Bad Air Quality



NAST<sup>219</sup>

The graphs illustrate the observed association between ground-level ozone (a component of smog) concentration and temperature in Atlanta and New York City (May to October 1988 to 1990) in parts per billion (ppb).219 The projected higher temperatures across the United States in this century are likely to increase the occurrence of high ozone concentrations, although this will also depend on emissions of ozone precursors and meteorological factors. Ground-level ozone can exacerbate respiratory diseases and cause short-term reductions in lung function.



Pollen production from ragweed grown in chambers at the carbon dioxide concentration of a century ago (about 280 parts per million [ppm]) was about 5 grams per plant; at today's approximate carbon dioxide level, it was about 10 grams; and at a level projected to occur about 2075 under the higher emissions scenario, 91 it was about 20 grams. 307


#### Working at the Local Level



Our weather (in South Carolina) is getting worse.

The changes in our weather are bad for us in a variety of ways.

The changes in our weather are happening because of the earth's changing climate.



#### The New Normal







# Conveying the Human Implications of Climate Change

A Climate Change Communication Primer for Public Health Professionals

Maibach E, Nisbet M, & Weathers M. (2011) Conveying the Human Implications of Climate Change: A Climate Change Communication Primer for Public Health Professionals. Fairfax, VA: George Mason University Center for Climate Change Communication.

#### Risks for Older Adults

Table I: Heal	Table I: Health Effects of Climate Change in the United States					
Weather Event	Health Effects	Populations Most Affected				
Heat Waves	Premature death  Heat-related illnesses such as heat stroke, heat exhaustion, and kidney stones	The elderly Children Diabetes Poor, urban residents People with respiratory diseases Those active outdoors (workers, athletes, etc.)				
Poor air quality	Increased asthma Increased chronic obstructive pulmonary disease (COPD) and other respiratory diseases	Children Those active outdoors (workers, athletes, etc.) The elderly People with respiratory diseases The poor				
Hurricanes	Death from drowning Injuries Mental health impacts such as depression and post- traumatic stress disorder Increased carbon monoxide poisoning Increased gastrointestinal illness Population displacement/homelessness	Coastal residents The poor The elderly Children				
Extreme rainfall and floods	Death from drowning Injuries Increased water-borne diseases from pathogens and water contamination from sewage overflows Increased food-borne disease	Residents in low-lying areas The elderly Children The poor Residents in the Southwestern U.S.				
Wildfires	Death from burns and smoke inhalation Injuries Eye and respiratory illness due to fire-related air pollution	People with respiratory diseases				
Droughts	Disruptions in food supply Changing patterns of crops, pests, and weed species Water shortages Malnutrition Food- and water-borne disease Emergence of new vector-borne and zoonotic disease	The poor The elderly Children				

heat waves
poor air quality
hurricanes
extreme rain/floods
drought

Maibach E, Nisbet M, & Weathers M. (2011) Conveying the Human Implications of Climate Change: A Climate Change Communication Primer for Public Health Professionals. Fairfax, VA: George Mason University Center for Climate Change Communication.


#### Risks for Older Adults



Towards a Quantitative Estimate of Future Heat Wave Mortality under Global Climate Change

Roger D. Peng, Jennifer F. Bobb, Claudia Tebaldi Larry McDaniel, Michelle L. Bell, and Francesca Dominici

doi: 10.1289/ehp.1002430 (available at http://dx.doi.org/)
Online 30 December 2010



#### Risks for Older Adults

# THE CLIMATE GAP

Inequalities in How Climate Change Hurts Americans & How to Close the Gap



http://dornsife.usc.edu/pere/documents/The\_Climate\_Gap\_Full\_Report\_FINAL.pdf

#### The New Normal

	Total Number of Households	Total Occupied Units	Black (not Hispanic)	Hispanic	Elderly (65 years or older)	Below Poverty Level
All Occupied Units	3,131,000	39.7%	58.5%	54.6%	37.5%	51.5%
Renters	1,608,900	48.1%	59.1%	58.4%	38.7%	56.3%
Homeowners	1,522,100	30.9%	57.4%	48.9%	36.8%	38.8%


Table 1. Percent of households without access to any air conditioning by race and SES — Los Angeles-Long Beach Metropolitan Area, California (2003)\*

Adapted from: American Housing Survey for the Los Angeles-Long Beach Metropolitan Area 2004 (USCB 2004).

The Climate Gap: Inequalities in How Climate Change Hurts Americans & How to Close the Gap

<sup>\*</sup> Percentages are likely an underestimate of the true value due to the fact that more than one category may apply to a single unit in the dataset.

#### The New Normal



The Climate Gap:
Inequalities in How
Climate Change Hurts
Americans & How to
Close the Gap

Figure 1. Percent change in mortality associated with 10°F increase in mean daily temperature by age group in nine California counties. May through September, 1999–2003 (Source: Basu and Ostro 2008).

## Seeing the Future

Climate is what you affect, weather is what gets you.

Myles Allen, Oxford University